arXiv:0909.1493v1 [math-ph] 8 Sep 2009

THE EXISTENCE AND UNIQUENESS OF SOLUTIONS TO N-BODY PROBLEM OF ELECTRODYNAMICS

VICTOR M. BOGDAN

Abstract. Given n charges interacting with each other according to Feynman's law. Let $(r_j(t), v_j(t))$ denote the position and velocity of the charge q_j. The list $y(t)$ of all such vectors is called a trajectory. A Lipschitzian trajectory $x(t), (t \leq 0)$, with continuous derivative, on which the velocities do not exceed some limiting velocity $v < c$, where c denotes the speed of light, is called an initial trajectory. A locally Lipschitzian trajectory $y(t)$ is called relativistically admissible if the velocities on it stay below the speed of light c. The author constructs operators Φ_j of a trajectory whose values $\Phi_j(y)(t)$ are linear transformations of R^3 into R^3. A point $t = t_1$ on a trajectory y is called singular if either some of the charges collide at the time t_1 or the determinant is zero for at least one of the transformations $\Phi_j(y)(t_1)$. The main result is the following: If $x(t)$ $(t \leq 0)$ is an initial trajectory with nonsingular point $t = 0$, then there exists a unique relativistically admissible trajectory $y(t)$, defined for t in an interval $I \subset \langle 0, \infty)$, extending the initial trajectory $x(t)$ and having the following properties. (1) No point t on the trajectory y is singular. (2) The trajectory represents a unique solution of the Newton-Einstein momentum-force system of equations under Lorentz forces induced by electromagnetic field in accord to Feynman's law for moving point charges. (3) The trajectory y represents the maximal global solution of the system.

Assume that we have a system of n charges interacting with each other according to Feynman's law for moving point chargesAssume that $y_j(t) = (r_j(t), v_j(t))$ denotes the position and velocity of the charge q_j. The list $y(t)$ of all such vectors is called a *trajectory* of the system. Let $\mathcal{D}$ denote the differential operator $\frac{d}{dt}$. Consider the Newton-Einstein system of equations with Lorentz forces

$$
\begin{aligned}
\mathcal{D}p_j &= q_j \sum_{k \neq j} (E_{jk} + \frac{1}{c}\, v_j \times [e_{jk} \times E_{jk}]) \;\; (j = 1, \ldots, n) \\
E_{jk} &= \frac{q_k}{4\pi\epsilon_0} \Big[\frac{e_{jk}}{|r_{jk}|^2} + \frac{|r_{jk}|}{c} \mathcal{D} \frac{e_{jk}}{|r_{jk}|^2} + \frac{1}{c^2} \mathcal{D}^2 e_{jk} \Big] \; (j \neq k)
\end{aligned}
\tag{1}
$$

where E_{jk} represents the intensity of the electric field acting onto the charge q_j and generated by the charge q_k, in accord with Feynman's law for moving point charges.

The quantity p_j denotes the relativistic momentum of the charge q_j. In the above formula r_{jk} represents the vector starting at a point on the trajectory of the charge q_k, when the wave was emitted at time t_{jk}, and ending at the point $r_j(t)$ at the

---

1991 *Mathematics Subject Classification.* 35L05, 53C50, 53C80, 78A25, 78A35, 81V10, 83C50.

Key words and phrases. nonlinear operators, Banach spaces, nonanticipating operators, differential equations, electrodynamics, n-body problem, electromagnetic fields, mathematical physics, relativity.

time t on the trajectory of the charge q_j, when the wave arrived. The vector e_{jk} denotes the unit vector in the direction of the vector r_{jk}.

A Lipschitzian trajectory $x(t)\,(t \le 0)$, with continuous derivative, on which the velocities do not exceed a certain limiting velocity v smaller than the speed of light c, is called an *initial trajectory.* A locally Lipschitzian trajectory $y(t)$ is called *relativistically admissible* if the velocities on it stay below the speed of light c, though they may approach c arbitrarily close.

Theorem 1. *Assume that the trajectory* $y(t)\,(t \ge 0)$, *extends an initial trajectory* $x(t)$ *and is relativistically admissible. Then for every such a trajectory* y *there exist unique functions*

$$t_{jk}(t) \text{ and } r_{jk}(t) \quad \text{for all} \quad t \in R$$

that appear in Feynman's formula (1) for the intensity of the electric field E_{jk}.

Define the operators of a trajectory y by the following formulas

$$\Phi_j = \Gamma(v_j) - [\frac{q_j q_k}{4\pi\epsilon_0 c^2 m_j}] \sum_{k \ne j} H_{jk} \circ G_{jk} \quad \text{for} \quad j = 1, 2, \ldots, n$$

where m_j is the relativistic mass of the charge q_j. The operators Γ, H_{jk}, and G_{jk} have their values in the space of linear transformations from the Euclidean space R^3 into R^3. They are given by the formulas for all $h \in R^3$

$$\begin{aligned} \Gamma(v_j)(h) &= h + \gamma_j^2 (u_j, h) u_j \\ H_{jk}(h) &= (h + \frac{1}{c}\, v_j \times \left[\, e_{jk} \times h \,\right]) \\ G_{jk}(h) &= |r_{jk}|^{-1} \{ h + [c - (e_{jk}, v_k(t_{jk}))]^{-1} (e_{jk}, h) [c e_{jk} + v_k(t_{jk})] \} \end{aligned}$$

where $u_j = \frac{1}{c} v_j$ and $\gamma_j = (1 - |v_j|^2 c^{-2})^{-1/2}$.

A point $t = t_1$ on a trajectory y is called *singular* if either some of the charges collide at the time t_1 or the determinant is zero for at least one of the transformations $\Phi_j(y)(t_1)$.

Theorem 2. *If* $x(t)\,(t \le 0)$ *is an initial trajectory with nonsingular point* $t = 0$, *then there exists a unique relativistically admissible trajectory* $y(t)$, *defined for* t *in an interval* $I \subset \langle 0, \infty)$, *extending the initial trajectory* $x(t)$ *and having the following properties.*

1. *No point* t *on the trajectory* y *is singular.*
2. *The trajectory* y *represents a solution of the Newton-Einstein system of equations with Lorentz forces induced by electromagnetic field in accord to Feynman's law for moving point charges (1).*
3. *The trajectory* y *cannot be extended any further preserving the above properties that is it represents the maximal global solution of the system (1).*

In order to prove the theorems the author makes use of the theory of generalized Lebesgue-Bochner-Stieltjes integral as developed in Bogdanowicz [2] and [3], the Banach contraction mapping theorem, and the works of Lorentz, Einstein, and Feynman.

Detailed proofs of the theorems will appear in *Quaestiones Mathematicae* [10].

References

[1] S. Bochner, *Integration von Funktionen, deren Werte die Elemente eines Vektorraumes sind*, Fund. Math. 20, (1933),p. 262–276.

[2] W.M. Bogdanowicz, (also known as V.M. Bogdan), *A Generalization of the Lebesgue-Bochner-Stieltjes Integral and a New Approach to the Theory of Integration*, Proc. of Nat. Acad. Sci. USA, Vol. 53, No. 3, (1965), p. 492–498. On the web at http://faculty.cua.edu/bogdan/rep/10.pdf

[3] ——— *An Approach to the Theory of Lebesgue-Bochner Measurable Functions and to the Theory of Measure*, Math. Annalen 164, (1966), p. 251–269. On the web at http://faculty.cua.edu/bogdan/rep/14.pdf

[4] ——— *Existence and uniqueness of extensions of volumes and the operation of completion of a volume.* Proceedings of the Japan Academy 42 (1966):571-576. On the web at http://faculty.cua.edu/bogdan/rep/15.pdf

[5] ——— *On volumes generating the same Lebesgue - Bochner integration.* Proceedings of the National Academy of Sciences USA 56 (1966):1399-1405. On the web at http://faculty.cua.edu/bogdan/rep/20.pdf

[6] W.M. Bogdanowicz and B. Kritt, *Radon - Nikodym differentiation of one vector-valued volume with respect to another.* Bulletin de l'Academie Polonaise des Sciences 15 (1967):479-486. On the web at
http://faculty.cua.edu/bogdan/rep/22.pdf

[7] V.M. Bogdan, *Existence and uniqueness of solutions for a class of non-linear–operator–differential equations arising in automatic spaceship navigation,* NASA, Scientific and Technical Information Branch, Technical Paper 1860, May (1981):1-18. On the web at http://faculty.cua.edu/bogdan/rep/58.pdf

[8] V.M. Bogdan, *Existence and uniqueness of solutions to nonlinear - operator - differential equations generalizing dynamical systems of automatic spaceship navigation.* Nonlinear Phenomena in Mathematical Sciences, Academic Press, (1982):123-136 On the web at http://faculty.cua.edu/bogdan/rep/60.pdf

[9] V.M. Bogdan, *Existence of solutions to differential equations of relativistic mechanics involving Lorentzian time delays,* Journal of Mathematical Analysis and Applications, Academic Press, 118, No. 2, September (1986),pp. 561-573. On the web at http://faculty.cua.edu/bogdan/rep/61.pdf

[10] V.M. Bogdan, *Feynman's Electromagnetic Fields Induced by Moving Charges and the Existence and Uniqueness of Solutions to N-Body Problem of Electrodynamics,* Quaestiones Mathematicae, vol. 32, (2009):1-87

[11] R. D. Driver and M. J. Norris, *A Collinear n-Body Problem of Classical Electrodynamics,* Nonlinear Phenomena in Mathematical Sciences, Vol. 4, pp. 329-334, Academic Press, New York, 1982.

[12] A. Einstein, *Zur Elektrodynamik bewegter Kōrper,* Annalen der Physik. 17, 891: 1905.
Enlish translation: *On the Electrodynamics of Moving Bodies,* The Principle of Relativity, Methuen and Company, 1923 and on the web at
http://www.fourmilab.ch/etexts/einstein/specrel/www/

[13] R. P. Feynman, *The Development of the Space-Time View of Quantum Electrodynamics,* Nobel Lecture, December 11, 1965. On the web at
http://nobelprize.org/nobel_prizes/physics/laureates/1965/feynman-lecture.html

[14] R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman Lectures on Physics* Vol. 1–3, Addison-Wesley, Reading, Mass., 1975.

[15] H. A. Lorentz, *Lectures on Theoretical Physics,* MacMillan& Co., London, 1927.

Department of Mathematics, McMahon Hall 207, CUA, Washington, D.C. 20064
E-mail address: `bogdan@cua.edu, web fttp://faculty.cua.edu/bogdan`